



## Alkyl deoxy-*arabino*-hexopyranosides: Synthesis, surface properties, and biological activities

Filipa V. M. Silva,<sup>a,b</sup> Margarida Goulart,<sup>a,b</sup> Jorge Justino,<sup>b</sup> Ana Neves,<sup>b</sup>  
Fernando Santos,<sup>c</sup> João Caio,<sup>a</sup> Susana Lucas,<sup>a</sup> Ana Newton,<sup>a</sup> Diana Sacoto,<sup>a</sup>  
Ester Barbosa,<sup>a</sup> Maria-Soledade Santos<sup>a</sup> and Amélia P. Rauter<sup>a,\*</sup>

<sup>a</sup>Centro de Química e Bioquímica, Departamento de Química e Bioquímica da Faculdade de Ciências da Universidade de Lisboa  
Edifício C8, 5º Piso, Campo Grande, 1749-016 Lisboa, Portugal

<sup>b</sup>Escola Superior Agrária de Santarém, Quinta do Galinheiro, 2001-904 Santarém, Portugal

<sup>c</sup>Centro de Ciências Moleculares e Materiais, Departamento de Química e Bioquímica,  
Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

Received 6 December 2007; revised 10 January 2008; accepted 11 January 2008

Available online 16 January 2008

**Abstract**—Octyl and dodecyl glycosides possessing 2-deoxy-*arabino*-hexopyranoside moieties belonging to the D- and L-series in their  $\alpha$ - and  $\beta$ -forms were synthesized by reaction of an acetyl protected glycal with octanol or dodecanol, catalyzed by triphenylphosphine hydrobromide, followed by deprotection. Their surface properties were studied and discussed in terms of the adsorption and aggregation parameters,  $pC_{20}$ , CMC, and  $\gamma_{CMC}$ . The antimicrobial activities were assessed using the paper disk diffusion and broth dilution methods. Both the octyl and dodecyl 2-deoxy  $\beta$ -D-glycosides inhibited significantly *Enterococcus faecalis*, a microbe also highly susceptible to dodecyl 2,6-dideoxy- $\alpha$ -L-*arabino*-hexopyranoside. This compound was particularly active against *Bacillus cereus* and *Bacillus subtilis*, presenting for both *Bacillus* species a minimal inhibitory concentration of the same order of magnitude and a minimal lethal concentration even smaller than that obtained for chloramphenicol, a bioactivity which remained unaltered after 1 year solution storage at 4 °C. In addition, activity over *Listeria monocytogenes* was also observed. Direct cytotoxicity and genotoxicity of the glycosides were determined by proliferative index (mitotic index) evaluation in peripheral human lymphocytes of healthy donors. All compounds induced acute toxicity effects, and the response was dose dependent for the  $\alpha$ -anomer of both the alkyl 2-deoxy-*arabino*-hexopyranosides and for the corresponding dodecyl  $\beta$ -anomer, what suggests that non-toxic but still bioactive concentrations may be found for these compounds.

© 2008 Elsevier Ltd. All rights reserved.

**Keywords:** Deoxy glycosides; Synthesis; Surface properties; Antimicrobial activity; Cytotoxicity.

\* Corresponding author. Tel.: +351 21 7500952; fax: +351 21 7500088; e-mail: [aprauter@fc.ul.pt](mailto:aprauter@fc.ul.pt)